



ELSEVIER

Journal of Computational and Applied Mathematics 114 (2000) 361–366

JOURNAL OF  
COMPUTATIONAL AND  
APPLIED MATHEMATICS

data, citation and similar papers at [core.ac.uk](http://core.ac.uk)

brought to

provided by Elsevier - F

# Discrete semi-classical orthogonal polynomials: generalized Charlier

M.N. Hounkonnou<sup>a,b,\*</sup>, C. Hounga<sup>a</sup>, A. Ronveaux<sup>c</sup>

<sup>a</sup>Unité de Recherche en Physique Théorique (URPT), Institut de Mathématiques et de Sciences Physiques (IMSP),  
B.P. 613 Porto-Novo, Benin

<sup>b</sup>Laboratoire de Physique Mathématique et de Mathématique Appliquée, Institut International de Recherche et de  
Formation (INIREF), B.P. 2740 Cotonou, Benin

<sup>c</sup>Facultés Universitaires Notre-Dame de la Paix, Rue de Bruxelles 61, 5000 Namur, Belgium

Received 10 October 1998; received in revised form 28 May 1999

## Abstract

Generalized Charlier polynomials are introduced as semi-classical orthogonal polynomials of class 1 with one parameter. Main characteristic data are established from the Laguerre–Freud equations generating the coefficients of the recurrence relation satisfied by the polynomials. © 2000 Elsevier Science B.V. All rights reserved.

MSC: 33C25; 39A70

Keywords: Orthogonal polynomial; Recurrence relation; Difference equation

## 1. *r*-generalized Charlier polynomials

Classical continuous and discrete orthogonal polynomials are defined by a positive weight  $\rho \equiv \rho(x)$ , solution of a Pearson differential or difference equation given, respectively, by

$$D(\sigma(x)\rho(x)) = \tau(x)\rho(x), \quad \Delta(\sigma(x)\rho(x)) = \tau(x)\rho(x), \quad (1.1)$$

where  $Df = df/dx = f'$ ,  $\Delta f(x) = f(x+1) - f(x)$ ;  $\sigma(x)$  is a polynomial of degree  $\leq 2$  and  $\tau(x)$  of degree 1. A striking difference between continuous and discrete case arises from the following considerations:

\* Corresponding author. Unité de Recherche en Physique Théorique (URPT), Institut de Mathématiques et de Sciences Physiques (IMSP), B.P. 613 Porto-Novo, Benin.

E-mail address: [hounkon@syfed.bj.refer.org](mailto:hounkon@syfed.bj.refer.org) (M.N. Hounkonnou)

Let us consider in both cases the equation satisfied by a new weight

$$\bar{\rho} \equiv \bar{\rho}(x) = [\rho(x)]^\lambda, \quad \lambda > 0.$$

In the continuous case, using differential equation (1.1)

$$\frac{\bar{\rho}'}{\bar{\rho}} = \lambda \frac{\rho'}{\rho} = \lambda \frac{\tau - \sigma'}{\sigma} = \frac{\lambda\tau - (\lambda - 1)\sigma' - \sigma'}{\sigma} \quad (1.2)$$

and the power of the classical weight is still a classical weight which satisfies:

$$(\bar{\sigma}\bar{\rho})' = \bar{\tau}\bar{\rho} \quad \text{with } \bar{\sigma} = \sigma \quad \text{and} \quad \bar{\tau} = \lambda\tau - (\lambda - 1)\sigma' \quad (1.3)$$

( $\bar{\tau}$  staying of first degree).

The discrete case is completely different. From (1.1)

$$\frac{\rho(x+1)}{\rho(x)} = \frac{\sigma(x) + \tau(x)}{\sigma(x+1)} \quad (1.4)$$

and now the  $\lambda$  power of this discrete weight becomes solution of

$$\frac{\bar{\rho}(x+1)}{\bar{\rho}(x)} = \left[ \frac{\sigma(x) + \tau(x)}{\sigma(x+1)} \right]^\lambda \quad (1.5)$$

which is never classical, and only semi-classical if  $\lambda$  is a positive integer. Classical Charlier polynomials correspond to  $\sigma(x)=x$  and  $\tau(x)=\mu-x$  with its orthogonal support being all integer  $i$ , ( $\mu > 0$ ). The solution of (1.4) is given by Nikiforov and Uvarov [4]

$$\rho(i) = \frac{\mu^i}{i!} \quad i = 0, 1, \dots \quad (1.6)$$

The  $r$ -generalized Charlier polynomials are defined from Eq. (1.1) with

$$\sigma(x) = x^r, \quad \tau(x) = \mu - x^r, \quad r \text{ is a positive integer}, \quad (1.7)$$

giving now for the weight  $\rho_r$

$$\frac{\rho_r(x+1)}{\rho_r(x)} = \frac{\mu}{(x+1)^r}, \quad (1.8)$$

the solution being

$$\rho_r(i) = \frac{\mu^i}{(i!)^r}, \quad i = 0, 1, \dots \quad (1.9)$$

For  $r=2$ , this weight is semi-classical of class  $s=1$  and the moments  $M_k(\mu) = \sum_{i=0}^{\infty} i^k \mu^i / (i!)^2$  can be expressed in terms of modified Bessel functions  $I_k$ ,  $k=0$  and  $1$

$$M_0(\mu) = I_0(2\sqrt{\mu}), \quad M_1(\mu) = \sqrt{\mu} I_1(2\sqrt{\mu}). \quad (1.10)$$

The recurrence relation for the moments easily obtained from Eq. (1.9) reads

$$M_{k+2}(\mu) = \mu M_k(\mu) - \sum_{i=1}^k \binom{k}{i} (-1)^i M_{k+2-i}(\mu). \quad (1.11)$$

The corresponding power of a Charlier weight (of parameter  $\sqrt{\mu}$ ) is given by

$$\rho_2^{(\mu)}(i) = [\rho^{(\sqrt{\mu})}(i)]^2, \quad i = 0, 1, \dots \quad (1.12)$$

These generalized Charlier polynomials with  $r = 2$  denoted  $\bar{C}_n^{(\mu)}(x)$  cannot be obtained from any weight modification of the Charlier polynomials  $C_n^\mu(x)$ , in contrast to the generalized Meixner case introduced in [5].

## 2. Recurrence coefficients for the generalized Charlier orthogonal polynomials ( $r = 2$ )

Let  $(P_n)_{n \geq 0}$  be a family of monic polynomials of degree exactly equal to  $n$  and orthogonal with respect to a positive weight  $\rho(x)$ . The three-term recurrence relation satisfied by the family  $(P_n)_{n \geq 0}$  writes as

$$\begin{aligned} P_{n+1}(x) &= (x - \beta_n)P_n(x) - \gamma_n P_{n-1}(x), \quad n \geq 1 \\ P_0(x) &= 1, \quad P_1(x) = x - \beta_0, \quad \gamma_n \neq 0 \quad \text{for all } n \geq 1. \end{aligned} \quad (2.1)$$

The family of orthogonal polynomials  $(P_n)_{n \geq 0}$  is said semi-classical of class  $s$ , with respect to a difference operator  $\Delta$  [2,3], if

$$\Delta(\sigma\rho) = \tau\rho, \quad (2.2)$$

$\tau(x)$  and  $\sigma(x)$  being given polynomials of fixed degree with

$$s = \max\{[\text{degree of } \tau(x)] - 1, [\text{degree of } \sigma(x)] - 2\}.$$

A crucial problem consists to deduce the recurrence coefficients  $\beta_n, \gamma_n$  as function of the coefficients contained in  $\tau(x)$  and  $\sigma(x)$  [1–3]. This leads to a coupled algebraic system of equations giving  $\beta_n$  and  $\gamma_n$  that are called Laguerre–Freud equations [2,3].

For

$$\tau(x) = \mu - x^2, \quad \sigma(x) = x^2, \quad (2.3)$$

the Laguerre–Freud equations for the generalized Charlier polynomials ( $r = 2$ ) reduce to

$$\gamma_{n+1} = \sum_{i=0}^{n-1} \beta_i + n\beta_n - \gamma_n - \beta_n^2 - \binom{n}{2} + \mu, \quad n \geq 1, \quad (\gamma_0 \equiv 0) \quad (2.4)$$

$$(\beta_{n+1} + \beta_n)\gamma_{n+1} = \sum_{i=0}^n \beta_i^2 + 2 \sum_{k=1}^n \gamma_k - n \sum_{i=0}^n \beta_i + (n+1)\gamma_{n+1} + \binom{n+1}{3}, \quad n > 1, \quad (2.5)$$

with the following initial values:

$$\beta_0 = \frac{M_1}{M_0} = \sqrt{\mu} \frac{I_1(2\sqrt{\mu})}{I_0(2\sqrt{\mu})}; \quad \gamma_1 = \mu - [\beta_0]^2, \quad (2.6)$$

$M_0$  and  $M_1$  being the moments of zero- and first-order, respectively. Then the coefficient  $\gamma_{n+1}$  can be uniquely expressed as function of  $\beta_n$  such that

$$\begin{aligned} \gamma_{n+1} &= n \sum_{i=0}^n (-1)^i \beta_{n-i} + 2 \sum_{i=1}^N i(\beta_{n+1-2i} - \beta_{n-2i}) \\ &\quad - \sum_{i=0}^n (-1)^i \beta_{n-i}^2 - \sum_{i=0}^{n-2} (-1)^i \binom{n-i}{2} + \mu - \beta_0^2 \end{aligned} \quad (2.7)$$

with  $N = \text{integer}[n/2, (n+1)/2]$ . The computation of  $\beta_n$  and  $\gamma_n$  values versus  $n$  for various  $\mu$ , using computer algebra, leads to the following remarks:

- (i) For each given  $\mu$ , the function  $\gamma_n$  versus  $n$  asymptotically tends to  $\mu$ , and this asymptotic value is reached as shorter as  $\mu$  is smaller; in particular for  $\mu \leq 0.1$ ,  $\gamma_n = \mu = \text{constant } \forall n$ ;
- (ii) For each given  $\mu$ , the function  $\beta_n$  versus  $n$  evolves to become equal to  $n$  and this equality occurs as more rapidly as smaller  $\mu$ ; in particular, for smaller  $\mu$  (typically for  $\mu \leq 0.1$ ), this function equals to  $n$  ( $\beta_n \equiv n$ )  $\forall n$ .

Whence, the three-term recurrence relation (2.1) satisfied by the family of the 2-generalized monic Charlier semi-classical orthogonal polynomials should be written as

$$P_{n+1}(x) = (x - n)P_n(x) - \mu P_{n-1}(x), \quad (2.8)$$

when  $n \rightarrow \infty$ .

Let mention that  $\beta_n$  and  $\gamma_n$  can be explicitly expressed in a representation involving exclusively the relevant parameters  $\mu$  and  $M_1$  and  $M_0$ . For example, the first coefficients, write as

$$\begin{aligned} \gamma_1 &= \mu - \beta_0^2, \\ \beta_1 &= \frac{\mu - \beta_0 + \beta_0^3}{\mu - \beta_0^2}, \\ \gamma_2 &= \frac{-\mu\beta_0(2\beta_0^2 + \beta_0 - 2\mu)}{(\beta_0^2 - \mu)^2}, \\ \beta_2 &= \frac{3\beta_0^5 + (2\mu + 1)\beta_0^4 - 3\mu\beta_0^3 - 4\mu^2\beta_0^2 + 2\mu^3}{\beta_0(\beta_0^2 - \mu)(2\beta_0^2 + \beta_0 - 2\mu)}. \end{aligned} \quad (2.9)$$

### 3. Structure relations and second-order difference equation

As any semi-classical orthogonal family,  $P_n(x)$  satisfies a structure relation which for the class  $s = 1$  reduces to

$$\sigma(x)\nabla P_n(x) = \sum_{i=n-2}^{n+2} \lambda_{n,i} P_i(x), \quad (3.1)$$

where  $\lambda_{n,i}$  are constants. These five coefficients can be computed [6] in general from the recurrence coefficients  $\gamma_n, \beta_n$ , and in the generalized Charlier case ( $r = 2$ ) reduce to ( $\lambda_{n,n+2} \equiv 0$ )

$$\lambda_{n,n+1} = n; \quad \lambda_{n,n} = -\binom{n}{2} + n\beta_n + \sum_{i=0}^{n-1} \beta_i; \quad \lambda_{n,n-1} = \gamma_n(\beta_{n-1} + \beta_n); \quad \lambda_{n,n-2} = \gamma_{n-1}\gamma_n. \quad (3.2)$$

Using basic recurrence (2.1), structure relation (3.1) can be reduced to

$$\sigma(x)\nabla P_n(x) = N_n(x)P_{n-1}(x) + M_n(x)P_n(x), \quad (3.3)$$

or, equivalently, to a similar equation using the following relation between  $\nabla$  and  $\Delta$ :

$$\nabla P_n(x) = \Delta P_n(x) - \Delta \nabla(P_n(x)) = \Delta P_n(x - 1), \quad (3.4)$$

where

$$\begin{aligned} M_n(x) &= \lambda_{n,n} - \frac{\lambda_{n,n-2}}{\gamma_{n-1}} + (x - \beta_n)\lambda_{n,n+1} \\ N_n(x) &= \lambda_{n,n-1} + \frac{\lambda_{n,n-2}}{\gamma_{n-1}}(x - \beta_{n-1}) - \lambda_{n,n+1}\gamma_n. \end{aligned} \quad (3.5)$$

In the same way, one can use another structure relation given by

$$(\sigma(x) + \tau(x))\Delta P_n(x) = \sum_{i=n-2}^{n-1} \tilde{\lambda}_{n,i} P_i(x) \quad (3.6)$$

such that

$$\alpha(x) = \sigma(x) + \tau(x) = \mu \quad (3.7)$$

and

$$\tilde{\lambda}_{n,n-1} = n\mu, \quad (3.8)$$

$$\tilde{\lambda}_{n,n-2} = \mu \left[ \binom{n}{2} - n\beta_{n-1} + \sum_{i=0}^{n-1} \beta_i \right]. \quad (3.9)$$

The structure relation (3.6) thus rewrites as

$$\mu\Delta P_n(x) = \tilde{N}_n(x)P_{n-1}(x) + \tilde{M}_n(x)P_n(x), \quad (3.10)$$

with

$$\begin{aligned} \tilde{M}_n(x) &= -\frac{\tilde{\lambda}_{n,n-2}}{\gamma_{n-1}} \\ \tilde{N}_n(x) &= \tilde{\lambda}_{n,n-1} + \frac{\tilde{\lambda}_{n,n-2}}{\gamma_{n-1}}(x - \beta_{n-1}). \end{aligned} \quad (3.11)$$

Using now Eqs. (3.3), (3.4) and (3.10), and after some trivial algebra consisting to eliminate  $P_{n-1}(x)$  from the equations, we obtain the following second-order difference equation:

$$\begin{aligned} & \left\{ \left[ \sum_{i=0}^{n-2} \beta_i + (1-n)\beta_{n-1} + \binom{n}{2} \right] x^3 + \left[ n\gamma_{n-1} + (n-1)\beta_{n-1}^2 \right. \right. \\ & \quad \left. \left. - \beta_{n-1} \left( \sum_{i=0}^{n-2} \beta_i + \binom{n}{2} \right) \right] x^2 \right\} \Delta \nabla P_n(x) \\ & + \left\{ \left[ (n-1)\beta_{n-1} - \sum_{i=0}^{n-2} \beta_i - \binom{n}{2} \right] x^3 + \left[ \beta_{n+1} \left( \sum_{i=0}^{n-2} \beta_i + \binom{n}{2} \right) \right. \right. \\ & \quad \left. \left. + (1-n)\beta_{n-1}^2 - n\gamma_{n-1} \right] x^2 + \gamma_{n-1}\gamma_n x + \gamma_{n-1}\gamma_n(\beta_n - n) \right\} \Delta P_n(x) \end{aligned}$$

$$\begin{aligned}
& + \left\{ \left[ n \sum_{i=0}^{n-2} \beta_i + (n - n^2) \beta_{n-1} + n \binom{n}{2} \right] x^2 \right. \\
& + \left[ \sum_{i=0}^{n-2} \beta_i^2 + 2 \sum_{0 \leq i < j \leq n-1} \beta_i \beta_j + (n-1)^2 \beta_{n-1}^2 - 2n \beta_{n-1} \sum_{i=0}^{n-2} \beta_i + n^2 \gamma_{n-1} - \binom{n}{2}^2 \right] x \\
& + \gamma_n \beta_n \left[ \sum_{i=0}^{n-2} \beta_i + (1-n) \beta_{n-1} + \binom{n}{2} \right] + \gamma_n \left[ \left( n^2 + \binom{n}{2} \right) \beta_{n-1} \right. \\
& \left. - n \left( \beta_{n-1}^2 + \gamma_{n-1} + \binom{n}{2} \right) \right] + \beta_{n-1}^2 \left[ n \left( \sum_{i=0}^{n-1} \beta_i - \binom{n}{2} \right) \right] \\
& \left. + \beta_{n-1} \left[ \gamma_n \sum_{i=0}^{n-1} \beta_i - \left( \sum_{i=0}^{n-1} \beta_i \right)^2 + \binom{n}{2}^2 \right] - n \binom{n}{2} \gamma_{n-1} \right\} P_n(x) = 0
\end{aligned}$$

satisfied by the 2-generalized Charlier polynomials.

## Acknowledgements

This work has been performed during the visit of M.N.H. at the Institut de Physique Théorique (Université Catholique de Louvain (Belgium)), supported by the Belgian Administration Générale pour la Coopération au Développement (AGCD). M.N.H. would like to thank Professor J.P. Antoine and all the staff of the institute for their kind hospitality. He also gratefully acknowledges Georges Destree of the Computer Center of the Université Libre de Bruxelles (Belgium) for numerous discussions and his positive contributions in the computational aspects of this work.

## References

- [1] S. Belmehdi, *Formes Linéaires et Polynômes Orthogonaux Semi-Classiques de classe  $s = 1$ . Description et Classification*, Thèse d'Etat, Université P. et M. Curie, Paris VI, 1990.
- [2] S. Belmehdi, A. Ronveaux, Laguerre–Freud's equations for the recurrence coefficients of semi-classical orthogonal polynomials, *J. Approx. Theory* 76 (1994) 351–368.
- [3] M. Foupouagnigni, M.N. Hounkonnou, A. Ronveaux, Laguerre–Freud equations for the recurrence coefficients of  $D_\omega$ -semi-classical orthogonal polynomials of class one, *J. Comput. App. Math.* 99 (1998) 143–154.
- [4] A.F. Nikiforov, V.B. Uvarov, *Special Functions of Mathematical Physics*, Birkhäuser, Basel, 1988.
- [5] A. Ronveaux, Discrete semi-classical orthogonal polynomials: generalized Meixner, *J. Approx. Theory* 46 (4) (1986) 403–407.
- [6] L. Salto Díaz, *Polinomios  $D_\omega$ -Semiclásicos*, Tesis Doctoral, Universidad de Alcalá de Henares, 1995.